



ES15 SPRAY & WIPE DISINFECTANT CLEANER (READY-TO-USE)

AVAILABLE IN USA / EPA 1839-220-64900

- Effective against a wide variety broad-spectrum of microorganisms including bacteria, antibiotic resistant bacteria and viruses

APPEARANCE	USA ODOR	pH-VALUE	SHELF LIFE
Blue Liquid	Mint	11-12	3 years

- Cleaner
- Disinfectant
- Deodorizer
- Sanitizer
- Bactericide
- Virucide

Alkyl (60% C14, 30% C16, 5% C12, 5% C18)
 dimethyl benzyl ammonium chloride..... 0.15% w/w
 Alkyl (68% C12, 32% C14)
 dimethyl ethylbenzyl ammonium chloride..... 0.15% w/w

- Food Processing Plants
- Food Service Establishments
- Hospitals/Healthcare settings
- Animal Housing Facilities/ Poultry & Swine Premises
- Washroom Facilities

Refer to label for complete instructions, dilutions, safety requirements and kill claims

Do not contaminate water, food, or feed by storage and disposal. Store in a dry place no lower in temperature than 10° C or higher than 50° C.

Rinse the emptied container thoroughly. Make the empty container unsuitable for further use. Dispose of the container in accordance with provincial requirements. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the Provincial Regulatory Agency or the Manufacturer.

Please refer to product label and SDS

Spray soiled area and wipe with a dry paper towel or lint free cloth.

DISINFECTING, DEODORIZING:

Remove gross filth or heavy soil prior to application of the product. Hold container 6-8 inches from the surface to be treated. Spray area until it is covered with the product. Allow surface to remain visibly wet for 5 minutes. No scrubbing necessary. Allow treated surfaces to air dry or wipe off with a clean cloth. When using on food contact surfaces: counter tops, stovetops, highchairs, kitchen appliances, thoroughly rinse all treated surfaces with potable water. This product must not be used to clean eating utensils, glassware and dishes.

is an effective bactericide and virucide in the presence of organic soil against:

Bactericidal Activity

Pseudomonas aeruginosa	2 Minutes
Salmonella enterica	2 Minutes
Staphylococcus aureus	2 Minutes
Multi Drug Resistant Acinetobacter baumannii Ceftazidime and Gentimicin	2 Minutes
Extended Spectrum Beta-Lactamase Escherichia coli	2 Minutes
Klebsiella pneumoniae New Dehli Metallo-Beta Lactamase Carbapenem Resistant	2 Minutes
Methicillin Resistant Staphylococcus aureus (MRSA)	2 Minutes
Vancomycin Resistant Enterococcus faecium	2 Minutes

Virucidal Activity

Human Immunodeficiency Virus Type-1 (HIV-1)	30 seconds
Avian Influenza Virus	30 seconds
Hepatitis B virus (HBV)	1 Minutes
Hepatitis C virus (HCV)	1 Minutes
Duck Hepatitis B Virus	1 Minutes
Bovine Viral Diarrhea Virus	1 Minutes
Human Rotavirus	1 Minutes
Feline Calicivirus	2 Minutes
Norovirus (Norwalk Virus)	2 Minutes
Canine Parvovirus	2 Minutes
Rhinovirus Type 14	2 Minutes
Rhinovirus Type 39	2 Minutes
Poliovirus Type 1	5 minutes



Mildewstat Activity

Aspergillus niger	10 minutes